



# Probability- Hints and Tips

## 1. Probability of Events:

### 1.1 Probability of one event:

- Probability of event E,  $P(E) = \frac{\text{Number of outcomes satisfying E}}{\text{Total number of outcomes}}$
- $0 \leq P(E) \leq 1$ , meaning that probabilities cannot be negative or greater than 1.
- $P(E) = 0$  means that event E can't happen.
- $P(E) = 1$  means event E is certain to happen.
- $P(E')$  is the probability that E does not happen.

### 1.2 Probability of multiple events:

- $P(A \cap B)$  is the probability that events A and B happen at the same time.
- $P(A' \cap B)$  is the probability that B happens but A does not happen at the same time
- $P(A \text{ or } B) = P(A) + P(B) - P(A \cap B)$ .

### 1.3 Conditional Probability:

- $P(A|B)$  is the probability that A happens given that B has happened.

### 1.4 Independent Events:

- Two events, A and B, are independent if the occurrence of one event does not affect the occurrence of the other e.g. flipping a coin and getting tails (A) and rolling a six-sided die and getting a 2 (B).
- For independent events,  $P(A \cap B) = P(A) * P(B)$ .
- Similarly for independent events:
  - $P(A|B) = P(A \cap B) / P(B) = P(A) * P(B) / P(B) = P(A)$
  - $P(A \cap B) = P(A) * P(B|A) = P(A) * P(B)$
  - $P(A \cap B) = P(B) * P(A|B) = P(B) * P(A)$

### 1.5 Expected Value:

- The expected value of a random variable is the sum of all the values that the random variable can take multiplied by the probability of the random variable being equal to that value.
- It is another word for the average value of the random variable.
- Formula is given by:  $E(X) = \sum x * P(X=x)$

## 2. Combinations and Permutations

- **Permutations:** The number of ways to arrange r objects from a set of n distinct objects.  $P(n,r) = \frac{n!}{(n-r)!}$ . Whenever the question mentions arrangements, think of permutations.
- **Combinations:** The number of ways to choose r objects from a set of n distinct objects, regardless of the order.  $C(n,r) = \frac{n!}{r!(n-r)!}$ . Whenever the question mentions choosing or choices, think of combinations.
- You need to be able to use your calculator to calculate different forms of permutations and combinations.
- To do this, first select the number n on your calculator, followed by the nCr button followed by the number r.



## Probability- Hints and Tips

- E.g. There are 11 students, how many choices of 5 a side teams can be made from those 11 students.
- In this question, the order does not matter, and we are looking for number of choices, so we are dealing with combinations.
- To calculate the number of combinations, we use our calculator and input 11, nCr, 5 and get an answer of 462.

### 3. Bernoulli Trials

#### 3.1 Bernoulli Trials

- A Bernoulli trial is one that only has two outcomes, success or failure.
- The trials are independent.
- Probability of success is p.
- Probability of failure is (1-p).

#### 3.2 Probability of k success in n Bernoulli Trials

- The formula for calculating the probability of r success in n events is given by the formula on page 33 of the log tables:

Dáilí dóchúlachta		Probability distributions
an dáileadh déthéarmach	$P(X = r) = \binom{n}{r} p^r q^{n-r}$ $r = 0 \dots n$	binomial distribution

- E.g. A fair die is thrown 5 times. Calculate the probability of getting 3 fours to 3 decimal places.
  - Probability of getting a four =  $\frac{1}{6}$
  - Probability of not getting a four =  $\frac{5}{6}$
  - Probability of 3 fours =  ${}_5C_3 * \left(\frac{1}{6}\right)^3 * \left(\frac{5}{6}\right)^2 = 0.032$

#### 3.3 Probability that the kth success occurs on the nth Bernoulli Trial

- In order for the kth success to happen on the nth trial, that means there must have been (k-1) success in (n-1) trials followed by a success on the nth trial.
- The formula is given by  ${}_{(n-1)}C_{(k-1)} * p^k * (1-p)^{(n-k)}$ .
- E.g. A card is randomly selected from a deck of cards and then placed back. Calculate the probability that the 3<sup>rd</sup> diamond is selected on the 10<sup>th</sup> card.
  - First calculate the probability of selecting 2 diamonds from the first 9 cards
  - ${}_9C_2 * \left(\frac{1}{4}\right)^2 * \left(\frac{3}{4}\right)^7 = 0.3$
  - Then multiply this by the probability of getting another diamond.
  - =  $0.3 * 0.25 = 0.075$ .